

Supporting Information

Organically Tempered Uranium(IV) Fluorooxalates with Layer Structures:

$(H_4TREN)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$ (TREN = tris(2-aminoethyl)amine) and

$(H_4APPIP)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$ (APPIP =

1,4-bis(3-amino-propyl)piperazine)

Chih-Min Wang,^{a,b} Yi-Ying Wu,^a Pei-Lin Chen,^a and Kwang-Hwa Lii^{a,b,*}

^aDepartment of Chemistry, National Central University, Chungli, Taiwan 320, R.O.C.

^bInstitute of Chemistry, Academia Sinica, Nankang, Taipei, Taiwan 115, R.O.C.

Figure S1. Experimental (top) and simulated (bottom) X-ray powder patterns for

$(H_4TREN)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$

Figure S2. The infrared spectrum of $(H_4TREN)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$ (KBr method).

Figure S3. Thermogravimetric analysis of $(H_4TREN)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$ in flowing air at 5 $^{\circ}C\ min^{-1}$.

Figure S4. Experimental (top) and simulated (bottom) X-ray powder patterns for

$(H_4APPIP)[U_2F_6(C_2O_4)_3] \cdot 4H_2O$.




Figure S1




Figure S2




Figure S3




Figure S4