Electronic Supplementary Information (ESI)

Reactions of a methylmercury zwitterionic thiolate complex [MeHg(Tab)]PF₆ with various ligands: relevance to methylmercury detoxification

Ai-Xia Zheng, a Hong-Xi Li, a Kai-Peng Hou, a Jing Shi, a Hui-Fang Wang, a Zhi-Gang Ren and Jian-Ping Lang a *a,b

^a College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou 215123, People's Republic of China

^b State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing 210093, P. R. China

Figure S1. The positive-ion ESI mass spectrum of [Hg(4,6-Me₂pymS)₂].

S3

Figure S2. The positive-ion ESI mass spectrum of 1.

S3

Figure S3. The positive-ion ESI mass spectrum of 2.

S3

Figure S4. The positive-ion ESI mass spectrum of 3.

S3

Figure S5. The positive-ion ESI mass spectrum of 4.

S4

Figure S6. The positive-ion ESI mass spectrum of 6.




Figure S1. The positive-ion ESI mass spectrum of [Hg(4,6-Me₂pymS)₂].




Figure S2. The positive-ion ESI mass spectrum of 1.




Figure S3. The positive-ion ESI mass spectrum of 2.




Figure S4. The positive-ion ESI mass spectrum of 3.




Figure S5. The positive-ion ESI mass spectrum of ${\bf 4}$.




Figure S6. The positive-ion ESI mass spectrum of 6.