

Supporting Information

Nitronyl and Imino Nitroxide Free Radicals as Precursors of Magnetic Phthalocyanine and Porphyrin Building Blocks

Emel Önal,^{a,b} Yusuf Yerli,^c Bunyemin Cosut,^a Guillaume Pilet,^b Vefa Ahsen,^a Dominique
Luneau*^b and Catherine Hirel*^a

^aGebze Institute of Technology, Department of Chemistry, 41400, Kocaeli, Turkey


E-mail: chirel@gyte.edu.tr; Fax ; +90 262 605 31 01 ; Tel : +90 262 605 30 21

^bUniversité Claude Bernard Lyon 1, Laboratoire des Multimatériaux et Interfaces (UMR 5615),
Campus de La Doua, 69622 Villeurbanne Cedex, France


E-mail : dominique.luneau@univ-lyon1.fr; Fax : +33 472 440 618 ; Tel : +33 472 431 418

^cGebze Institute of Technology, Department of Physics, 41400 Kocaeli, Turkey


a)




b)




c)




d)



e)



f)



g)




Figure S1: X-band ESR spectra of mono radicals (a) **2** , (b) **3** , (c) **4** , (d) **5** , (e) **7** , (f) **8** , (g) **9** in CHCl_3 ($c = 10^{-4}$ M) at room temperature.





Figure S2: ESI mass spectrum of nitronyl nitroxide **3**.