


Supporting Information

Conjugated Energetic Salts Based on 3,3'-((1*E*,1'*E*)-(2,4,6-trinitro-1,3-phenylene)bis(ethene-2,1-diyl))bis(2,4,6-trinitrophenol)


Junhui Xu,^[a] Jianping Wei,^[a] Fangmei Li,^[a] Qingguo Ma ^[a] and Xinhua Peng*^[a]

School of Chemical Engineering, Nanjing University of Science and Technology, Nanjing 210094, China. E-mail: xhpeng@mail.njust.edu.cn Fax: (+) 86-025-84317639; Tel: (+) 86-025-84315520

Supporting Information





Comment 1
Comment 2



Bruker Daltonics flexAnalysis


printed: 2013/4/7 11:04:04

Supporting Information




Supporting Information



201300918_13 26 (0.689) Cm (22:28)



1: TOF MS ES-
2.16e6




Supporting Information




Supporting Information

201300918_22_26 (0.680) Cm (26:27)

1: TOF MS ES-
2.96e6




Supporting Information




(D₂O exchange result for Ammonium salt **4a**)



Supporting Information




Supporting Information





Supporting Information





Supporting Information





Supporting Information




Supporting Information





Supporting Information





Supporting Information




Supporting Information





Supporting Information





Supporting Information





Supporting Information




Supporting Information





Supporting Information




Supporting Information



Supporting Information



Supporting Information



Supporting Information

