

A supramolecular polymer with dual-responsiveness constructed by crown ether based molecular recognition

Danyu Xia and Min Xue



Department of Chemistry, Zhejiang University, Hangzhou 310027, P. R. China Fax: +86-571-8795-3189; Tel: +86-571-8795-3189;
E-mail: xuemin@zju.edu.cn

Electronic Supplementary Information (10 pages)

1. *Synthesis of monomer 1*

S2

*1. Synthesis of monomer **I***






Figure S2. ^{13}C NMR spectrum (CDCl_3 , 293 K, 125 MHz) of **4**.




Figure S3. Electrospray ionization mass spectrum of **4**. Main peak: m/z 705.8 $[\text{M} + \text{Cl}]^-$ (100%).




Figure S4. ^1H NMR spectrum (CDCl_3 , 293 K, 400 MHz) of **5**.




Figure S5. ^{13}C NMR spectrum (CDCl_3 , 293 K, 125 MHz) of **5**.





Figure S6. Electrospray ionization mass spectrum of **5**. Main peak: m/z 540.4 [M + H]⁺ (100%).






Figure S8. ^{13}C NMR spectrum (DMSO, 293 K, 125 MHz) of **1**.




Figure S9. Electrospray ionization mass spectrum of **1**. Main peak: m/z 540.3 $[\text{M} - \text{PF}_6]^+$ (100%).