

Supporting Information

Figures S-1 and S-2 show details regarding the transverse and longitudinal relaxivities of *MBICs* and *MBIClusters* as a function of hydrodynamic size. This material is available free of charge via the Internet at <http://pubs.acs.org>.




Figure S-1. Transverse relaxation rates of *MBICs* and *MBIClusters* with different intensity average diameters with a) 3.5K-6.8K H₂N-PEO-PAA, and b) 3.5K-9.5K H₂N-PEO-PAA as a function of iron concentration. Non-crosslinked *MBICs* are represented by the smallest of the average particle diameters in each case.




Figure S-2. Longitudinal relaxation rates of MBICs and *MBIClusters* with different intensity average diameters with a) 3.5K-6.8K H₂N-PEO-PAA, and b) 3.5K-9.5K H₂N-PEO-PAA as a function of iron concentration. Non-crosslinked MBICs are represented by the smallest of the average particle diameters in each case.




Figure S-3. ¹H NMR spectrum of *tboc*-NH-PEO-*b*-PtBA in CDCl₃ (top) and H₂N-PEO-*b*-PAA in D₆-DMSO (bottom)




Figure S-4. SEC analysis of *tboc*-NH-PEO-*b*-PtBA

Corresponding Author

*Judy Riffle, jriffle@vt.edu or judyriffle@aol.com